

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## CISTACEÆ.

- HELIANTHEMUM, Tourn.—II. Canadense, Michx.; common.
- HUDSONIA, L.—II. cricoides, L.; Tottenville, Staten Island, Le Roy.—II. tomentosa, Nutt.; abundant along the sea shore, May, June.
- LECHEA, L.—L. major, Michx.; common.—L. thymifolia, Pursh; not uncommon, particularly on the south side of L. I.—L. Novæ-Cæsareæ, Austin; not uncommon; abundant at Tottenville, Staten Island, W. H. L.—L. minor, Lam.; common with numerous variations.

#### DROSERACEÆ.

DROSERA, L.—D. rotundifolia, L.; common.—D. longifolia, L.; New Jersey, Austin: common in swamps on Long Island, Coles, Allen.

#### HYPERICACEÆ.

- ASCYRUM, L.—A. stans, Michx.; Long Island, Le Roy.—A. Crux-Andrew, L.; South side of Staten Island; South Amboy; Bergen Point, N. J., many years ago I used to find it in the neighborhood of 56th St. and the Hudson River, W. H. L.
- HYPERICUM, L.—II. prolificum, L.; Var. densiflorum; South Amboy, T. F. A.—II. adpressum, Barton; New Jersey, Austin.—II. angulosum, Michx.; Cedar swamps at Weehawken, Torr. Cat., but not seen of late.—II. perforatum, L.; very common.—II. corymbosum, Muhl; common.—II. mutilum, L.; very common.—II. Canadense, L.; very common; Var. major; Rockaway, L. I., W. H. L.—II. Sarothra, Michx.; very common.
- ELODES, Adans.—E. Virginica, Nutt.; common.

## ELATINACEÆ.

ELATINE, L.—E. Americana, Arnott; Banks of the Passaic; Ponds, L. I., T. F. A.; Westchester Co.; Mead, in State Flora; (Hudson R. at Peekskill, W. H. L.).

#### CARYOPHYLLACEÆ.

- DIANTHUS, L.—D. Armeria, L.; common.
- SAPONARIA, L.—S. officinalis, L.; common.
- VACCARIA, Medik.—V. vulgaris, Host.; Newtown Station of the Flushing R. R., Ruger.
- SILENE, L.—S. stellata, Ait.; common.—S. inflata, Smith; hill-sides on Staten Island, near the Narrows, T: F. A: Spuyten Duyvel, Westchester Co., W. H. L.; New York City, Austin.—S. Penn sylvanica, Michx.; common; still flourishes in its native habitat in Central Park.—S. Virginica, L.; reported, but no station named.—S. Armeria, L.; New Jersey, Austin; occasional by road sides, W. H. L.—S. antirrhina, L.; common.—S. noctiflora; cultivated grounds and road sides, Flushing etc., W. H. L.; Inwood, W. W. Denslow.
- LYCHNIS, Tourn.—L. vespertina, Sibth.; in similar situations with Silene noctiflora.—L. Githago, Lam.; grain fields, common.

- ARENARIA, L.—A. serpyllifolia, L.; very common.—A. squarrosa, Michx.; sandy soil, N. J., T. F. A.—A. lateriflora, L.; not uncommon.—A. perloides, L.; sea shore, Staten Island, Long Island.
- STELLARIA, L.—S. media, Smith; very common.—S. longifolia, Muhl.; New Jersey, Austin, W. H. L.; bogs and overflowed woods, Torr. Cat.; north of Astoria, Ruger.
- CERASTIUM, L.—C. velgetum, L.; shady rocks along N. R. R. of N. J. above junction with Erie R. R., T. F. Allen; New Jersey, very rare, Austin.—C. viscosum, L.; very common.—C. nutans, Raf.; woods, N. J., Torr. Cat., Austin.—C. oblongifolium, Torr.; abundant on the hills of Staten Island, not in Torr. Cat.; Palisades, N. J.; Austin, Le Roy.—C. arvense, L.; Fort Lee, T. F. Allen.
- SAGINA, L.—S. procumbens, L.; Austin; near Hempstead, W. H. L.; "Borders of creeks and springs, and in rather moist sandy soils; on Long Island abundant, also near the city of New York." State Flora.—S. apetala, L.; Austin; dry hill sides, near Manhattanville, State Flora.
- SPERGULARIA, Pers.—S. rubra, Presl., var. compestris; abundant along the New Jersey shores.—S. salina, Presl.; Coney Island, F. J. Bumstead; Maspeth and Canarsie, M. Ruger.—S. media, Presl.; common on New York and Long Islands, W. H. L.; Var. macretarpa, I have a well marked specimen of this plant, handed me some years since, while yet fresh, by Prof. Jas. Hyatt, who gathered it, as I understood, at Hoboken. W. H. L.
- SPERGULA, L.—S. arvensis, L.; rather common in cultivated grounds; rare in New Jersey, Austin.

ANYCHIA, Michx.—A. dichotoma, Michx.; common.

SCLERANTHUS, L.—S. annuns, L.; very common.

MOLLUGO, L.—M. verticillata, L.; very common.

# PORTULACACEÆ.

PORTULACA, Tourn.—P. oleracea, L.; common. CLAYTONIA, L.—C. Virginica, L.; common.

# MALVACEÆ.

- ALTHÆA, L.—A. officinalis; not infrequent along the north shore of Long Island; Newtown, T. F. A.; Flushing, State Flora; Glen Cove, Coles; etc.
- MALVA, L.—M. rotundifolia, L.; common.—M. sylvestris, L.; Westchester Co., Dr. Mead, in State Flora.—M. moschata, L.; a frequent escape.
- SIEA, L.—S. spinosa, L.; "Sandy hills, N. J." Eddy in Torr. Cat. Le Roy.
- ABUTILON, Tourn.—A. Avicennæ, Gærtn.; not uncommon.
- KONTELETZKYA, Presl.—K. Virginica, Presl.; "Salt marshes near Oyster Bay." Mitchill in Torr. Cat. Hackensack meadows, near "Pen Horn City" on Erie R. R., T. F. Allen.
- HIBISCUS, L.—II. Moscheutes, L.; common.